

Chama marina. Vitre. Dinn. Blaehlyen.

Sunt Concha marinae equivalves, aqua- vel in aquilatero tenuis vel crassiss testa, laxes, vel transversim striata triquetra, rotabiliter umbo natum, et oblique incurvata, ventri- colix, cervic, seu rostro umbo crassior, et utroque cuiuslibet valva late in cardine at- tingente, illaque incurvatura est lateri carinis fossilium conditorum in uno latere lon- giorum, in altero vero breviorum efformantes; superficies omnium interna laxis ex- ultit, externa autem varia est, prout ex descriptione specimen sub sequentium pat- bie.

Chama marina laxes tenuis testa.

Chama marina laxis tenuis testa maior albicans fascia lati fulci circa in- am oram cincta tripliciter representata

I. è latere fossula cordi-
formis longioris.

I. è parte umonis.

III. è latere fossula cor-
diformis brevioris.



Tradem medicocris obscure albicans fascia
pluribus fulci, et minus, seu dilute sultis in-
sionis. An Chama lavis Rond. sij. Blaet
Dinn. Blaehlyen. Gein. 154. Concha lavis Rond.
Bon. 47. et sequentibus. Chama lavis Glae-
meris Indica Rumph. t. 45. l. G.

Eius interna facies cum articulatione
apparente, et vestigio adhaesionis ani-
malis, glabrente, levigata, in qua ob-
splendorum coloris in interna superficie
clarior apparent, quam externe in glo-
bo naturali sita.

