

*METHODI
correctæ, et auctæ.
Iconibusque illustrata.*

TESTACEA MARINA

in suas debitas Clases. Genera, et Species distribuendi
secundum suas notas characteristicas

PARS TERTIA

Cochleæ marinae breviores, et magis contractæ
idest

Cochleæ marinae ore, et turbine
parum, aut nihil elongatis.

qua commode in tres classes dividuntur, nimis rūm:
in Triangulares, Globosæ, et Orbiculares.

Triangulares

et tribus rectis vel quasi rectis lineis constant,
qua simul unita tres angulos constituant
recti lineos, vel tantillum incurvatos,
aut quasi rectilineos:

et sic Trochos marinos efformant,
vel Cochleas marinas Trochiformes.

Sie Meer-Höppi, oder Höppi-formige Meer-Schnäcken,
quibus annumeranda est Cochlea calata.

Der ausgestochene Meer-Schnügel,
cum inter has, et levigentes medium quasi teneat.

Globosæ,

qua præter turbinem tantillum elongatum, ac ventricosum
os quoque notabiliter eminens habent,
sicque ad globi figuram quodammodo accedunt.
ad has præcipue pertinent.

Cochleæ terrestres, et marinæ,
figurâ sua ad eas proximè accedentes,

et propterea Terrestriformes dictæ
Eis-Schnäcken, und Eis-Schnäcken-formige Meer-Schnäcken,
qua iterum sunt simplices vel umbilicatae.

medium inter has obtinere videntur
Echinophoræ, Neritæ, et Cochlea marina Neritoidæ,
cum Calcaribus marinæset Cochlea laciniata
Meer-Höl, Neritische, und Neritoidesische Meer-Schnäcken,
mit den Meer-Höppen sc.
qua aut Turbinem tantillum elongatum habent,
aut os notabiliter eminens.



11b